

Localization and Pattern Formation in BBGKY Hierarchy

Antonina N. Fedorova, Michael G. Zeitlin

Mathematical Methods in Mechanics Group
Institute of Problems of Mechanical Engineering (IPME RAS)
Russian Academy of Sciences
Russia, 199178, St. Petersburg, V.O., Bolshoj pr., 61
zeitlin@math.ipme.ru, anton@math.ipme.ru
<http://www.ipme.ru/zeitlin.html>
<http://www.ipme.nw.ru/zeitlin.html>

Abstract

A fast and efficient numerical-analytical approach is proposed for modeling complex behaviour in the BBGKY hierarchy of kinetic equations. Numerical modeling shows the creation of various internal structures from localized modes, which are related to the localized or chaotic type of behaviour and the corresponding patterns (waveletons) formation.

Presented: GAMM Meeting, 2004, Dresden, Germany.

Published: Proc. Appl. Math. Mech. (PAMM), 4, 564, Wiley-VCH, 2004.

Localization and Pattern Formation in BBGKY Hierarchy

Antonina N. Fedorova and Michael G. Zeitlin*

IPME RAS, St. Petersburg, V.O. Bolshoj pr., 61, 199178, Russia

A fast and efficient numerical-analytical approach is proposed for modeling complex behaviour in the BBGKY hierarchy of kinetic equations. Numerical modeling shows the creation of various internal structures from localized modes, which are related to the localized or chaotic type of behaviour and the corresponding patterns (waveletons) formation.

Copyright line will be provided by the publisher

We consider the application of a new numerical/analytical technique based on local nonlinear harmonic analysis approach (multiscale decomposition) for the description of complex (non-equilibrium) behaviour of the classical statistical ensembles, considered in the framework of the general BBGKY hierarchy (kinetics equations). We restrict ourselves to the rational/polynomial type of nonlinearities (with respect to the set of all dynamical variables) that allows to use our results from [1],[2], which are based on the so called multiresolution framework [3] and the variational formulation of initial nonlinear (pseudodifferential) problems. Our main goals are an attempt of classification and construction of a possible zoo of nontrivial (meta) stable states: high-localized (nonlinear) eigenmodes, complex (chaotic-like or entangled) patterns, localized (stable) patterns (waveletons). The last case is a good image for plasma modeling in fusion state (energy confinement). It should be noted that the class of smoothness (related at least to the appearance of chaotic/fractal-like type of behaviour) of the proper functional space under consideration plays a key role in the following [2]. Let M be the phase space of an ensemble of N particles ($\dim M = 6N$) with coordinates $x_i = (q_i, p_i)$, $i = 1, \dots, N$, $q_i = (q_i^1, q_i^2, q_i^3) \in R^3$, $p_i = (p_i^1, p_i^2, p_i^3) \in R^3$, $q = (q_1, \dots, q_N) \in R^{3N}$. Individual and collective measures are: $\mu_i = dx_i = dq_i dp_i$, $\mu = \prod_{i=1}^N \mu_i$. Our constructions can be applied to the following general Hamiltonians: $H_N = \sum_{i=1}^N \left(\frac{p_i^2}{2m} + U_i(q) \right) + \sum_{1 \leq i \leq j \leq N} U_{ij}(q_i, q_j)$, where the potentials $U_i(q) = U_i(q_1, \dots, q_N)$ and $U_{ij}(q_i, q_j)$ are restricted to rational functions of the coordinates. Let L_s and L_{ij} be the Liouvillean operators (vector fields) and $F_N(x_1, \dots, x_N; t)$ be the hierarchy of N -particle distribution function, satisfying the standard BBGKY-hierarchy (V is the volume):

$$\frac{\partial F_s}{\partial t} + L_s F_s = \frac{1}{V^s} \int d\mu_{s+1} \sum_{i=1}^s L_{i,s+1} F_{s+1} \quad (1)$$

Our key point in the following consideration is the proper nonperturbative generalization of the perturbative multiscale approach of Bogolyubov. The infinite hierarchy of distribution functions satisfying system (1) in the thermodynamical limit is: $F = \{F_0, F_1(x_1; t), \dots, F_N(x_1, \dots, x_N; t), \dots\}$, where $F_p(x_1, \dots, x_p; t) \in H^p$, $H^0 = R$, $H^p = L^2(R^{6p})$ (or any different proper functional space), $F \in H^\infty = H^0 \oplus H^1 \oplus \dots \oplus H^p \oplus \dots$ with the natural Fock space like norm (guaranteeing the positivity of the full measure):

$$(F, F) = F_0^2 + \sum_i \int F_i^2(x_1, \dots, x_i; t) \prod_{\ell=1}^i \mu_\ell. \quad (2)$$

First of all we consider $F = F(t)$ as a function of time only, $F \in L^2(R)$, via multiresolution decomposition which naturally and efficiently introduces the infinite sequence of the underlying hidden scales. Because the affine group of translations and dilations generates multiresolution approach, this method resembles the action of a microscope. We consider a multiresolution decomposition of $L^2(R)$ [3] (of course, we may consider any different and proper for some particular case functional space) which is a sequence of increasing closed subspaces $V_j \in L^2(R)$ (subspaces for modes with fixed dilation value): $\dots V_{-2} \subset V_{-1} \subset V_0 \subset V_1 \subset V_2 \subset \dots$. The closed subspace $V_j (j \in \mathbf{Z})$ corresponds to the level j of resolution, or to the scale j and satisfies the following properties: let W_j be the orthonormal complement of V_j with respect to V_{j+1} : $V_{j+1} = V_j \oplus W_j$. Then we have the following decomposition: $\{F(t)\} = \bigoplus_{-\infty < j < \infty} W_j =$

$V_c \bigoplus_{j=0}^{\infty} W_j$ in case when V_c is the coarsest scale of resolution. The subgroup of translations generates a basis for the

fixed scale number: $\text{span}_{k \in \mathbf{Z}} \{2^{j/2} \Psi(2^j t - k)\} = W_j$. The whole basis is generated by action of the full affine group: $\text{span}_{k \in \mathbf{Z}, j \in \mathbf{Z}} \{2^{j/2} \Psi(2^j t - k)\} = \text{span}_{k, j \in \mathbf{Z}} \{\Psi_{j,k}\} = \{F(t)\}$. In multidimensional case we may consider polynomial tensor bases, e.g. in $n = 2$ case we may use the rectangle lattice of scales and one-dimensional wavelet decomposition:

* Corresponding author: e-mail: zeitlin@math.ipme.ru, <http://www.ipme.ru/zeitlin.html>, <http://www.ipme.nw.ru/zeitlin.html>

$f(x_1, x_2) = \sum_{i,\ell,j,k} \langle f, \Psi_{i,\ell} \otimes \Psi_{j,k} \rangle \Psi_{i,\ell} \otimes \Psi_{j,k}(x_1, x_2)$, where the basis functions $\Psi_{i,\ell} \otimes \Psi_{j,k}$ depend on two scales 2^{-i} and 2^{-j} . We obtain our multiscale/multiresolution representations (formulae (3) below) via the variational wavelet approach for the following formal representation of the BBGKY system (1) (or its finite-dimensional nonlinear approximation for the n -particle distribution functions) with the corresponding obvious constraints on the distribution functions. Let L be an arbitrary (non)linear differential/integral operator with matrix dimension d (finite or infinite), which acts on some set of functions from $L^2(\Omega^{\otimes n})$: $\Psi \equiv \Psi(t, x_1, x_2, \dots) = (\Psi^1(t, x_1, x_2, \dots), \dots, \Psi^d(t, x_1, x_2, \dots))$, $x_i \in \Omega \subset \mathbf{R}^6$, n is the number of particles: $L\Psi \equiv L(Q, t, x_i)\Psi(t, x_i) = 0$, $Q \equiv Q_{d_0, d_1, d_2, \dots}(t, x_1, x_2, \dots, \partial/\partial t, \partial/\partial x_1, \partial/\partial x_2, \dots, \int \mu_k) = \sum_{i_0, i_1, i_2, \dots=1}^{d_0, d_1, d_2, \dots} q_{i_0 i_1 i_2 \dots}(t, x_1, x_2, \dots) \left(\frac{\partial}{\partial t}\right)^{i_0} \left(\frac{\partial}{\partial x_1}\right)^{i_1} \left(\frac{\partial}{\partial x_2}\right)^{i_2} \dots \int \mu_k$. Let us consider the N mode approximation for the solution as the following ansatz: $\Psi^N(t, x_1, x_2, \dots) = \sum_{i_0, i_1, i_2, \dots=1}^N a_{i_0 i_1 i_2 \dots} A_{i_0} \otimes B_{i_1} \otimes C_{i_2} \dots(t, x_1, x_2, \dots)$. We shall determine the expansion coefficients from the following conditions: $\ell_{k_0, k_1, k_2, \dots}^N \equiv \int (L\Psi^N) A_{k_0}(t) B_{k_1}(x_1) C_{k_2}(x_2) dt dx_1 dx_2 \dots = 0$. Thus, we have exactly dN^n algebraical equations for dN^n unknowns $a_{i_0, i_1, \dots}$. This variational approach reduces the initial problem to the problem of solution of functional equations at the first stage and some algebraical problems at the second. The solution is parametrized by the solutions of two sets of reduced algebraical problems, one is linear or nonlinear (depending on the structure of the operator L) and the rest are linear problems related to the computation of the coefficients of the algebraic equations. The solution of the equations (1) has the following multiscale decomposition via high-localized eigenmodes

$$F(t, x_1, x_2, \dots) = \sum_{(i,j) \in \mathbb{Z}^2} a_{ij} U^i \otimes V^j(t, x_1, \dots), \quad (3)$$

$$V^j(t) = V_N^{j, \text{slow}}(t) + \sum_{l \geq N} V_l^j(\omega_l t), \quad \omega_l \sim 2^l, \quad U^i(x_s) = U_M^{i, \text{slow}}(x_s) + \sum_{m \geq M} U_m^i(k_m^s x_s), \quad k_m^s \sim 2^m,$$

which corresponds to the full multiresolution expansion in all underlying time/space scales. The formulae (3) give the expansion into a slow and fast oscillating parts. So, we may move from the coarse scales of resolution to the finest ones for obtaining more detailed information about the dynamical process. In this way one obtains contributions to the full solution from each scale of resolution or each time/space scale or from each nonlinear eigenmode. It should be noted that such representations give the best possible localization properties in the corresponding (phase)space/time coordinates. Formulae (3) do not use perturbation techniques or linearization procedures. Numerical calculations are based on compactly supported wavelets and related wavelet families [3] and on evaluation of the accuracy on the level N of the corresponding cut-off of the full system (1) regarding norm (2): $\|F^{N+1} - F^N\| \leq \varepsilon$. Fig. 1 demonstrates the appearance of localized (meta) stable pattern (wavelet), which can be considered, e.g., as a model for fusion state in plasma.




Fig. 1 Localized wavelet pattern.

References

- [1] A.N. Fedorova and M.G. Zeitlin, in: *Quantum Aspects of Beam Physics*, Ed. P. Chen, (World Scientific, Singapore, 2002), p. 527, p. 539; arXiv preprints: physics/0101006, 0101007, quant-ph/0306197, 0405044; 0406009; 0406010; J. Mod. Optics, **51**, 1105 (2004).
- [2] A.N. Fedorova and M.G. Zeitlin, in: *Progress in Nonequilibrium Green's Functions II*, Ed. M. Bonitz, (World Scientific, Singapore, 2003), p. 481; arXiv preprint: physics/0212066, physics/0212115.
- [3] Y. Meyer, *Wavelets and Operators* (Cambridge Univ. Press, 1990).